



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : G01N 33/53, 53/567, C07K 16/00	A1	(11) International Publication Number: WO 99/42830 (43) International Publication Date: 26 August 1999 (26.08.99)
(21) International Application Number: PCT/US99/03170 (22) International Filing Date: 12 February 1999 (12.02.99) (30) Priority Data: 60/075,504 23 February 1998 (23.02.98) US 09/111,938 8 July 1998 (08.07.98) US 09/175,504 20 October 1998 (20.10.98) US (71) Applicant (for all designated States except US): DIADEXUS LLC [US/US]; 3303 Octavius Drive, Santa Clara, CA 95054 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BURCZAK, John [US/US]; Apartment 301, 1526 Village Club, Santa Clara, CA 95054 (US). WILKINSON, Francis, E. [US/US]; 3838 Jarvis Avenue, San Jose, CA 95118 (US). (74) Agents: LICATA, Jane, Massey et al.; Law Offices of Jane Massey Licata, 66 E. Main Street, Marlton, NJ 08053 (US).		(81) Designated States: CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: METHODS USING PLA ₂ AS A MARKER OF METASTASES AND FOR THE DIAGNOSIS OF SELECTED CANCERS (57) Abstract <p>The present invention provides methods of monitoring a cancer in a patient which has not metastasized for the onset of metastasis by measuring PLA₂ levels in the patient. Also provided are methods for diagnosing a cancer in a patient and monitoring progression, remission, response to therapy and stabilization of a cancer in a patient based upon PLA₂ levels.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

METHODS USING PLA₂ AS A MARKER OF METASTASES AND FOR THE DIAGNOSIS OF SELECTED CANCERS

INTRODUCTION

5 This application is a continuation of application
Serial No. 09/175,504, filed October 20, 1998 which is a
continuation-in-part of application Serial No. 09/111,938,
filed July 8, 1998, pending, which is a continuation-in-part
of provisional application Serial No. 60/075,504, filed
10 February 23, 1998.

FIELD OF THE INVENTION

 This invention relates to use of PLA₂ as a metastatic
marker for monitoring cancers which have not metastasized for
the onset of metastasis and for diagnosing metastatic cancers.
15 PLA₂ is demonstrated herein to be a metastatic marker for a
number of different metastatic cancers including prostate,
breast, colorectal, ovarian and testicular cancer. This
invention also relates to methods of monitoring selected
cancers including breast, ovarian and testicular cancer for
20 progression, remission, response to therapy and stabilization
in patients by monitoring PLA₂ levels in these patients.
Further, the invention relates to methods which aide in
identification of cancers, including testicular and ovarian
cancer, among individuals who have not yet been diagnosed with
25 these cancers by detection of elevated PLA₂ levels.

- 2 -

BACKGROUND OF THE INVENTION

Extracellular Phospholipase A₂ (PLA₂) enzymes appear to mediate a variety of responses including cellular proliferation, chemotaxis and inflammation. There are two major groups of extracellular PLA₂ enzymes: pancreatic or group I; and rheumatoid arthritis synovial fluid (RASF) or group II. The group I enzymes function in digestion and in modulating proliferation and chemotaxis. Currently, RASF-PLA₂ is predominantly believed to play a role in inflammatory responses including arthritis, septic shock and lung injury. The level of RASF-PLA₂ is regulated at the mRNA level by a variety of agents including interleukin-6, interleukin-1 and tumor necrosis factor, all of which are involved in inflammatory responses. As used hereinbelow "PLA₂" refers to group II or RASF-PLA₂. PLA₂ includes native PLA₂ (whole or a breakdown product), a native complex of molecules including PLA₂, or native chemically modified PLA₂.

Membrane-associated phospholipase A₂, M-PLA₂, has been disclosed as one of five factors suspected to play a role in the tumor cell processes and metastasis in human breast cancer (Yamashita et al., *Surgery*, 1995, 117, 601-608). Further, higher M-PLA₂ levels have been reported in breast cancer patients with distant metastasis and in patients with scirrhous carcinoma as compared to other histological types (Yamashita, *Cancer*, 1994, 69, 1166-1170). PLA₂ has also been reported to be expressed exclusively in gastric cancer cells with a low grade of differentiation and appears to be intensified in the invading zone of the tumor (Murata et al., *Br. J. Cancer*, 1993, 68(1), 103-111). Elevated M-PLA₂ was also detected in three gastric cancer cell lines (MKN28, KATO III and AZ521), in a pancreatic cancer cell line (SUIT-2), in a colonic cancer cell line (SW1116) and a hepatoblastoma cancer cell line (HuH-6). (Yamashita et al., *Clinica Chimica Acta*, 1994, 228, 91-99). In addition, patients with T2-T4 tumors or stage II-IV cancers of the lung, breast and

- 3 -

digestive organs had significantly higher levels of M-PLA₂ in comparison to stage I and T1 cancers.

PLA₂ levels have also been reported to be increased in peritoneal and pleural effusions in patients with various
5 types of cancers including gastric, breast, pancreatic, bile duct, lung, liver, esophageal and uterine cancer and cirrhotic ascites (Abe et al., *Int. J. Cancer (Pred. Oncol.)*, 1997, 74 245-250).

In addition, U.S. Patent 5,747,264 discloses a
10 diagnostic assay for detection of PLA₂ protein or PLA₂ mRNA in cells, tissues or body fluids which can be used to detect the presence of cancers, and in particular, prostate cancer. This method of quantifying PLA₂ protein levels is particularly useful for discriminating benign prostatic hyperplasia and
15 prostate cancer.

It has now been found that levels of PLA₂ in a patient can be used to diagnose and monitor various cancers for the onset of metastasis. It has also been found that determining the amount of PLA₂ in a bodily fluid of a patient
20 is useful in diagnosing various cancers and in monitoring progression, remission, response to therapy and stabilization of certain cancers in patients.

SUMMARY OF THE INVENTION

Toward these ends, and others, it is an object of the
25 present invention to provide a method of monitoring various cancers which have not metastasized for the onset of metastasis which comprises identifying a patient having a cancer that is not known to have metastasized and measuring an amount of PLA₂ in a biological fluid obtained from the
30 patient, wherein elevated levels of PLA₂ in the sample are indicative of metastasis. As demonstrated herein, PLA₂ is a metastatic marker for cancers, including prostate, breast, colorectal, ovarian and testicular cancer.

- 4 -

It is another object of the present invention to provide a method of diagnosing a metastatic cancer in a patient which comprises obtaining a sample of biological fluid from a patient; and detecting the level of PLA_2 in a sample of
5 biological fluid, wherein elevated levels of PLA_2 in the sample are indicative of metastatic cancers.

It is another object of the present invention to provide a method for monitoring cancers, including breast, ovarian and testicular cancer, for progression, remission,
10 response to therapy and stabilization in patients suffering from such cancers which comprises measuring levels of PLA_2 in biological fluid samples obtained from the patient at selected times, wherein an increase in the measured levels of PLA_2 over time is indicative of progressive cancer, a decrease in the
15 measured levels of PLA_2 over time is indicative of remission or response to therapy of the cancer, and no change in the measured levels of PLA_2 over time is indicative of stabilization of the cancer.

It is another object of the present invention to
20 provide a method to aide in identification of ovarian or testicular cancer, among individuals who have not yet been diagnosed with such cancers which comprises detecting elevated PLA_2 levels in a biological fluid obtained from a patient suspected of suffering from cancer.

25 Other objects, features, advantages and aspects of the present invention will become apparent to those of skill in the art from the following description. It should be understood, however, that the following description and the specific examples, while indicating preferred embodiments of
30 the invention, are given by way of illustration only. Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following description and from reading the other parts of the present disclosure.

- 5 -

DESCRIPTION OF THE INVENTION

Procedures used for detecting, diagnosing, monitoring, staging, and prognosticating cancer are of critical importance to the outcome of patients suffering from cancer. Cancer patients are closely monitored following initial therapy and during adjuvant therapy to determine response to therapy and to detect persistent or recurrent disease or metastasis. Another important step in managing cancer is to determine the stage of the patient's disease. Stage determination has potential prognostic value and provides criteria for designing optimal therapy. Generally, pathological staging of cancer is preferable over clinical staging because the former gives a more accurate prognosis. However, clinical staging would be preferred were it at least as accurate as pathological staging because it does not depend on an invasive procedure to obtain tissue for pathological evaluation. Staging of cancer would be improved by detecting new markers in cells, tissues or bodily fluids which could differentiate between different stages of invasion. Accordingly, there is a need for cancer markers which are more sensitive and specific in detecting cancer recurrence and for increasingly sensitive and accurate methods for the staging of a cancer in humans to determine whether or not such cancer has metastasized and for monitoring the progress of a cancer in a human which has not metastasized for the onset of metastasis.

For example, cancer of the prostate is the most prevalent malignancy in adult males, excluding skin cancer, and is an increasingly prevalent health problem in the United States. In 1996 in the United States alone, it was estimated that 41,400 deaths would result from this disease, indicating that prostate cancer is second only to lung cancer as the most common cause of death in the same population. If diagnosed and treated early, when the cancer is still confined to the prostate, however, the chances of cure are significantly

- 6 -

higher. Further, treatment decisions for an individual are linked to the stage of prostate cancer present in that individual. A common classification of the spread of prostate cancer has been established by the American Urological Association (AUA). The AUA divides prostate tumors into four stages, A to D. Stage A, microscopic cancer within prostate, is further subdivided into stages A1 and A2. Sub-stage A1 is a well-differentiated cancer confined to one site within the prostate. Treatment generally includes observation, radical prostatectomy, or radiation. Sub-stage A2 is a moderately to poorly differentiated cancer at multiple sites within the prostate. Treatment generally includes radical prostatectomy or radiation. Stage B, which is characterized by a palpable lump within the prostate, is also subdivided into stages B1 and B2. In sub-stage B1 the cancer forms a small nodule in one lobe of the prostate. In sub-stage B2 the cancer forms large or multiple nodules, or occurs in both lobes of the prostate. Treatment for both sub-stages B1 and B2 involves either radical prostatectomy or radiation. Stage C is a large cancer mass involving most or all of the prostate and is also further subdivided into two stages. Sub-stage C1 is characterized by the cancer forming a continuous mass that may extend beyond the prostate. Sub-stage C2 is characterized by the cancer forming a continuous mass that invades the surrounding tissue. Treatment for both sub-stages C1 and C2 is radiation with or without drugs to address the cancer. The fourth stage, Stage D is metastatic cancer and is also subdivided into two stages. Sub-stage D1 is characterized by the cancer appearing in the lymph nodes of the pelvis. Sub-stage D2 is characterized by the cancer appearing in tissues beyond the lymph nodes. Treatment for both sub-stages D1 and D2 is systemic drugs to address the cancer as well as pain.

However, current prostate cancer staging methods are limited. Existing methods for diagnosing prostate cancer such as prostatic specific antigen (PSA), digital examination, and

- 7 -

transurethral ultrasound tests have difficulty discriminating between prostate cancer stages A, B, C, and D. In fact, the existing PSA diagnostic tests detect 20-28% of patients with benign prostatic hyperplasia (BPH) and 62-61% of prostate cancer patients with PSA blood levels above approximately 99% of the normal population. In addition, as many as 50% of prostate cancers initially staged as A2, B, or C have metastasized and are actually stage D. However, accurate identification of metastasis is important because patients with metastatic cancers have a poorer prognosis and require significantly different therapy than those with localized cancers.

An incorrect initial diagnosis also occurs in up to 25% of patients with testicular tumors. However, the lifetime probability of developing testicular cancer is 0.2% for an American white male. The cause of testicular cancer is unknown, however, it is associated with both congenital and acquired factors. From a treatment standpoint, testicular cancer is divided into two major categories, nonseminomas and seminomas. In the commonly used staging system for nonseminomas, a stage A lesion is confined to the testis; in stage B there is regional lymph node involvement in the retroperitoneum; and in stage C, there is distant metastasis. For seminomas, a stage I lesion is confined to the testis; in stage II, the lesion has spread to the retroperitoneal lymph nodes; and in stage III, the lesion has supradiaphragmatic nodal or visceral involvement.

The most common symptom of testicular cancer is painless enlargement of the testis. Acute testicular pain resulting from intertesticular hemorrhage occurs in about 10% of cases. Patients are often asymptomatic upon presentation but about 10% may exhibit back pain, cough or lower extremity edema. A testicular mass or diffuse enlargement of the testis can be detected by physical examination in most cases. Several biochemical markers are used for the diagnosis and

- 8 -

treatment of this disease including human chorionic gonadotropin (hCG), alpha-fetoprotein and LDH. However, as evidenced by the high rate of initial misdiagnosis, better markers are required.

5 The most widely used biomarker for epithelial ovarian cancer, CA125, has also been reported to lack sensitivity and specificity (Xu et al. *JAMA*, 1998, 280 (8), 719-723). In 1995, an estimated 26,600 new cases of ovarian cancer were diagnosed in the United States. Approximately one in 70 women
10 will develop ovarian cancer during her lifetime. An estimated 14,500 deaths in 1995 resulted from ovarian cancer. It causes more deaths than any other cancer of the female reproductive system.

 Ovarian cancer often does not cause any noticeable
15 symptoms. Some possible warning signals, however, are an enlarged abdomen due to an accumulation of fluid or vague digestive disturbances (discomfort, gas or distention) in women over 40; rarely there will be abnormal vaginal bleeding. Periodic, complete pelvic examinations are important; a Pap
20 test does not detect ovarian cancer. Annual pelvic exams are recommended for women over 40.

 As with most cancers, the risk of ovarian cancer increases with age. The rate is highest among women over 60 years of age. Women who have never had children are twice as
25 likely to develop ovarian cancer as women who have. Women who already have been diagnosed with breast, intestinal or rectal cancers appear to be at increased risk of developing ovarian cancer. Early age of first pregnancy, early menopause and the use of oral contraceptives appear to reduce the risk of
30 ovarian cancer.

 Surgical treatment usually involves the removal of one or both ovaries, the uterus and the fallopian tubes. If the cancer is detected early, especially in younger women, it is possible that only the cancerous ovary will be removed.
35 Radiation and chemotherapy options also are available to

- 9 -

prevent or slow the recurrence of the ovarian cancer or the spread of cancer to other parts of the body following surgery.

It has now been found that PLA_2 levels are significantly higher in human patients with metastatic cancer
5 as compared to cancer which has not yet metastasized.

For example, it has now been found that PLA_2 is significantly higher in human patients with metastatic Stage D prostate cancer as compared to prostate cancer which has not yet metastasized, i.e cancer in Stage A, B, or C. PLA_2 levels
10 have also been demonstrated to be elevated in patients suffering from other progressive metastatic cancers such as breast cancer, colorectal cancer, testicular cancer and ovarian cancer. Accordingly, the present invention relates to methods of diagnosing metastatic cancer and monitoring
15 cancers which have not metastasized for the onset of metastasis.

In this method, a patient with cancer not known to have metastasized is identified. This is accomplished by a variety of means known to those of skill in the art. For
20 example, in the case of prostate cancer, patients are typically diagnosed with prostate cancer following digital rectal examination, serum levels of prostate specific antigen (PSA), transrectal ultrasound, and/or needle biopsy of the prostate and surrounding tissue. For ovarian cancer, patients
25 are typically diagnosed with ovarian cancer following pelvic examination. CA125 is the most widely biomarker for detection and management of epithelial ovarian cancer. Recently, plasma lysophosphatidic acid (LPA) levels have been suggested as a potential biomarker for ovarian cancer and other gynecological
30 cancers (Xu et al. *JAMA*, 1998, 280 (8), 719-723). In the case of testicular cancer, patients are typically diagnosed with testicular cancer following physical examination or detection of biochemical markers including human chorionic gonadotropin (hCG), alpha-fetoprotein and LDH. A sample of bodily fluid
35 is then obtained from this patient. Bodily fluids useful in

- 10 -

the present invention include blood, urine, saliva, or any other bodily secretion or derivative thereof. By blood it is meant to include whole blood, plasma, serum, or any derivative of blood, preferably serum. PLA₂ protein or mRNA levels in
5 the sample of bodily fluid are then determined.

Assay techniques that can be used to measure PLA₂ protein or mRNA encoding PLA₂ in a bodily fluid sample derived from a patient having cancer are well-known to those of skill in the art. Such assay methods include, without limitation,
10 radioimmunoassays, reverse transcriptase PCR (RT-PCR) assays, immunohistochemistry assays, *in situ* hybridization assays, competitive-binding assays, Western Blot analyses, ELISA assays and proteomic approaches: two-dimensional gel electrophoresis (2D electrophoresis) and non-gel based
15 approaches such as mass spectrometry or protein interaction profiling. Among these, enzyme immunoassays, and in particular enzyme linked immunosorbent assays (ELISAs) are frequently preferred to detect a gene's expressed protein in biological fluids. An ELISA assay initially comprises
20 preparing an antibody, if not readily available from a commercial source, specific to PLA₂, preferably a monoclonal antibody. In addition, a reporter antibody generally is prepared which binds specifically to PLA₂. The reporter antibody is attached to a detectable reagent such as a
25 radioactive, fluorescent or enzymatic marker, for example horseradish peroxidase enzyme or alkaline phosphatase. To carry out the ELISA, antibody specific to PLA₂ is incubated on a solid support, e.g., a polystyrene dish, that binds the antibody. Any free protein binding sites on the dish are then
30 covered by incubating with a non-specific protein such as bovine serum albumin (BSA). Next, the bodily fluid sample to be analyzed is incubated in the dish, during which time PLA₂ binds to the specific antibody attached to the polystyrene dish. Unbound sample is washed out with buffer. A reporter
35 antibody specifically directed to PLA₂ and linked to a

- 11 -

detectable reagent, i.e. alkaline phosphatase, is placed in the dish resulting in binding of the reporter antibody to the bound PLA₂. Unattached reporter antibody is then washed out and bound reporter antibody is detected. For example, if the
5 detectable reagent is alkaline phosphatase, reagents for alkaline phosphatase activity, including a colorimetric substrate are added to the dish. Immobilized alkaline phosphatase, linked to PLA₂ antibodies, produces a colored reaction product. The amount of color developed in a given
10 time period is proportional to the amount of PLA₂ protein present in the sample. As will be obvious to those of skill in the art upon this disclosure, other reporter antibodies and means for detecting these antibodies can also be used. Quantitative results typically are obtained by reference to
15 a standard curve.

A competition assay can also be used wherein antibodies specific to PLA₂ are attached to a solid support and labeled PLA₂ and sample obtained from the patient having cancer are passed over the solid support. The amount of label
20 detected attached to the solid support can then be correlated to a quantity of PLA₂ in the sample.

Nucleic acid methods can also be used to detect PLA₂ mRNA in a bodily fluid obtained from a patient having cancer. Examples of nucleic acid methods include, but are not limited
25 to polymerase chain reaction (PCR), ligase chain reaction (LCR) and nucleic acid based amplification (NASABA). Reverse transcriptase PCR (RT-PCR) also provides a powerful tool useful in the detection of the presence of a specific mRNA population in a complex mixture of thousands of other mRNA
30 species.

Of the proteomic approaches, 2D electrophoresis is a technique well known to those in the art. Isolation of individual proteins from a sample such as serum is accomplished using sequential separation of proteins by
35 different characteristics usually on polyacrylamide gels.

- 12 -

First, proteins are separated by size using an electric current. The current acts uniformly on all proteins, so smaller proteins move farther on the gel than larger proteins. The second dimension applies a current perpendicular to the first and separates proteins not on the basis of size but on the specific electric charge carried by each protein. Since no two proteins with different sequences are identical on the basis of both size and charge, the result of a 2D separation is a square gel in which each protein occupies a unique spot. Analysis of the spots with chemical or antibody probes, or subsequent protein microsequencing can reveal the relative abundance of a given protein and the identity of the proteins in the sample.

Additional methods obvious to those of skill in the art upon this disclosure for determining PLA₂ levels in bodily fluids can also be used.

Without limiting the instant invention, typically, for a quantitative diagnostic enzyme immunoassay, a positive result indicating that the cancer in the patient being tested or monitored has metastasized is one in which bodily fluid levels of the cancer marker, PLA₂, are elevated above an established enzyme immunoassay (EIA) cut-off of 4.5 ng/ml. However, as will be obvious to those of skill in the art upon this disclosure, alternative EIA cutoffs may be established depending upon the acceptable number of false positive or false negative results for a particular patient group. In addition, in another embodiment, elevated levels of PLA₂ can be determined by comparing measured amounts in the sample of bodily fluid obtained from the human patient having cancer with amounts of this marker in the same bodily fluid type of normal human controls or previously measured amounts in the same patient. That is, if PLA₂ in serum is being measured in the patient, this amount is compared with the amount of PLA₂ in serum of normal human controls or the amount of PLA₂ measured in serum of the same patient previously. By "normal

- 13 -

human control" it is meant a random grouping of males who have not been diagnosed with prostate cancer or any other type of cancer. An increase or elevation in PLA₂ in the patient versus the amount in normal human controls or the amount previously measured in the same patient is associated with a cancer which has metastasized.

Table 1 shows the percentage of individuals identified as positive based upon measurement of PLA₂ in serum by an ELISA with an EIA cut-off of 4.5 ng/ml in normal human controls, patients having benign prostatic hyperplasia, patients having prostatitis, patients having untreated prostate cancer in Stage A, patients having untreated prostate cancer in Stage B, patients having untreated prostate cancer in Stage C, and patient newly diagnosed and treated for progressive metastasis from prostate cancer in Stage D.

Table 1
Serum Levels of PSA and PLA₂

Groups	N	PLA ₂ Positive (%) (cutoff 4.5)	PSA Positive (%)
Random Males	1573	3.4	0.0
BPH	70	2.9	27.1
Prostatitis	20	15.0	30.0
Prostate Cancer Stage A	14	7.1	78.6
Prostate Cancer Stage B	41	7.3	87.8
Prostate Cancer Stage C	17	11.8	88.2
Prostate Cancer Stage D	58	81.0	87.9

The PSA Positive value for Random Males is based upon random PSA testing. As demonstrated in this table, the percent of

- 14 -

individuals positive for PLA₂ increases significantly in patients with Stage D prostate cancer as compared to either normal human controls or patients with prostate cancer in Stage A, B or C. Accordingly, measurement of PLA₂ provides a useful means for diagnosing metastatic cancer and/or monitoring the onset of metastasis in patients with prostate cancer which has not yet metastasized.

Table 2 shows the percentage of individuals identified as positive based upon measurement of PLA₂ in serum by an ELISA with an EIA cut-off of 4.5 ng/ml in patients having breast, colorectal, ovarian or testicular cancer classified as either localized, progressive metastatic, metastatic in remission or metastatic stable.

Table 2
Serum Levels of PLA₂

Cancer Type	Localized		Progressive Met		Met Remission		Met Stable	
	N	% Pos	N	% Pos	N	% Pos	N	% Pos
Breast	4	25.0	24	83.3	8	75.0	9	44.4
Colorectal	3	33.3	20	75.0	7	85.7	5	80.0
Ovarian	1	0.0	36	72.2	1	0.0	3	33.3
Testicular	6	50.0	14	64.3	14	28.6	7	28.6

As demonstrated in this table, the percent of individuals positive for PLA₂ also increases significantly in patients with progressive breast, colorectal, ovarian and testicular metastatic cancers as compared to patients with localized cancers. Accordingly, measurement of PLA₂ provides a useful means for monitoring the onset of metastasis in patients with various types of cancer, including prostate, breast, colorectal, ovarian and testicular cancer. Further, measurement of serum PLA₂ levels at the time of diagnosis of

- 15 -

prostate, breast, colorectal, ovarian or testicular cancer or subsequently thereafter serves as an aide not only in the diagnosis of the cancer but also in determining if the newly diagnosed cancer is metastatic.

- 5 Further, as shown in Table 2, the percent of individuals positive for PLA₂ decreases in patients with breast, ovarian or testicular metastatic cancer that is in remission or stabilized, from the elevated percent of individuals positive for PLA₂ with progressive metastasis.
- 10 Accordingly, measurement of PLA₂ provides a useful means to monitor progression, remission, response to therapy or stabilization of various cancers over time. Levels of PLA₂ can be determined in biological fluid samples obtained from the patient at selected times. Times for determining PLA₂
- 15 levels for monitoring the progression, remission, response to therapy or stabilization of cancer in a patient can be routinely selected by one of skill in the art in accordance with the patient's history and the type of cancer which is being monitored. An increase in the measured levels of PLA₂
- 20 in a patient over time is indicative of progressive cancer. A decrease in the measured levels of PLA₂ in a patient over time is indicative of remission or response to therapy of the cancer. No change in the measured levels of PLA₂ in a patient over time is indicative of stabilization of the cancer.
- 25 As also demonstrated in Table 2, elevated PLA₂ levels in a bodily fluid obtained from a patient are also indicative of breast, colorectal, ovarian or testicular cancer in the patient. The percent of individuals positive for PLA₂ with breast, colorectal, ovarian or testicular cancer that is
- 30 localized, progressive metastatic, metastatic in remission, or metastatic stable is generally greater than random healthy males or females.

Patients diagnosed early with cancer generally have a much greater five-year survival rate as compared to the

35 survival rate for patients diagnosed with distant metastasized

- 16 -

cancer. Thus, diagnostic methods which are more sensitive and early detection of cancers are clearly needed. Accordingly, the present invention also relates to methods for diagnosing cancers, and in particular ovarian cancer or testicular cancer, in a patient by measuring PLA₂ in bodily fluids of the patient suspected of suffering from cancer. In these methods, levels of PLA₂ at least two standard deviations above levels of PLA₂ determined in healthy males are indicative of ovarian or testicular cancer. Levels of PLA₂ in patients suspected of suffering from ovarian or testicular cancer are determined in a biological fluid obtained from the patient in accordance with well known assays as described herein.

Other objects, features, advantages and aspects of the present invention will become apparent to those of skill in the art based upon this description. It should be understood, however, that the description, while indicating preferred embodiments of the invention, is given by way of illustration only. Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the present disclosure.

The following nonlimiting examples are provided to further illustrate the present invention.

EXAMPLES

25 **Example 1: Quantitative immunoassay for Type II PLA₂**

Microtiter plates were prepared for assays by coating a purified monoclonal antibody (SK088-3C6.16.2; Roshak et al., *J. Biol. Chem.*, 1994, 269, 25999-26005) against type II PLA₂ at a concentration of 2.0 µg/ml in 10 mM Tris-HCl buffer at pH 8.0 overnight at 2-8°C. The microtiter wells were then washed four times with 10 mM Tris-HCl at pH 7.5 containing 150 mM NaCl and 0.05% Tween-20 (TBS-T) and blocked with 1.0% BSA (Sigma A-7888) in 50 mM Tris-HCl at pH 8.0 for one hour at 21-25°C (room temperature). The microtiter plates were then

- 17 -

washed four times again with TBS-T before immediate use in the assay.

For the immunocapture assay, 200 μ l of 0.5 μ g/ml of a second monoclonal antibody (SK097-1E8.5.2; Roshak et al.,
5 *J. Biol. Chem.*, 1994, 269, 25999-26005) against type II PLA₂ conjugated to biotin (Hinatownich et al., *J. Nucl. Med.*, 1987, 28, 1294-1302) in sample diluent (SPD) buffer (50 mM Tris-HCl at pH 8.0, 300 mM NaCl, 0.1% BSA, 0.05% Tween-20, 1% mouse serum (Jackson ImmunoResearch Labs 015-000-120), and 0.02%
10 NaN₃) was added to each well. Also added to the microtiter wells at this step was 10 μ l of either type II PLA₂ standard (Levin et al. *Protein Expr. Purif.* 1992, 3, 27-35) in fetal calf serum (HyClone A-1111-L) or patient serum. The plates were then incubated for 1 hour at 21-25°C with shaking. After
15 the incubation, the mixture was removed, the microtiter plates were washed 4 times with TBS-T and 100 μ l of avidin-alkaline phosphatase (Sigma A-2527) at 1.0 μ g/ml in SPD buffer were added. After a one hour incubation at 21-25°C with shaking, the wells were washed again four times with TBS-T. Finally
20 200 μ l of p-nitrophenyl phosphate at 1.0 mg/ml in 10 mM diethanolamine at pH 9.7 were added to the wells and incubated for 30 minutes at 21-25°C with shaking. Assay absorbance at 405 nm was read for each microtiter well. Assay absorbance values for purified type II PLA₂ standards were used to
25 generate a standard curve, and absorbance values for patient sample assays were compared to the standard curve to determine type II PLA₂ concentrations.

Example 2: Quantitative Assay of Prostate Specific Antigen (PSA) in Serum

30 PSA levels in serum samples of the same patients were quantified using the TANDEM-E PSA ImmunoEnzyMetric Assay (Hybritech, Inc. Sand Diego, CA) in accordance with the manufacturer's instructions.

- 18 -

What is claimed is:

1. A method of monitoring a cancer in a patient which has not metastasized for the onset of metastasis comprising:

5 (a) identifying a patient having a cancer that is not known to have metastasized;

(b) periodically analyzing a sample of bodily fluid from the patient for PLA₂; and

(c) detecting levels of PLA₂ in the sample of
10 bodily fluid obtained from the patient, wherein elevated levels of PLA₂ in a sample of bodily fluid are indicative of a cancer which has metastasized.

2. The method of claim 1 wherein the bodily fluid is serum.

15 3. The method of claim 1 wherein the analysis of the bodily fluid for PLA₂ is by ELISA.

4. The method of claim 3 wherein the established enzyme immunoassay cutoff is 4.5 ng/ml.

20 5. The method of claim 1 wherein the cancer is breast, colorectal, ovarian, prostate or testicular cancer.

6. A method of diagnosing a metastatic cancer in a patient comprising:

(a) obtaining a sample of biological fluid from a patient; and

25 (b) detecting the level of PLA₂ in a sample of biological fluid, wherein elevated levels of PLA₂ in the sample are indicative of a metastatic cancer.

7. The method of claim 6 wherein the bodily fluid is serum.

- 19 -

8. The method of claim 6 wherein the analysis of the bodily fluid for PLA₂ is by ELISA.

9. The method of claim 8 wherein the established enzyme immunoassay cutoff is 4.5 ng/ml.

5 10. A method of diagnosing ovarian or testicular cancer in a patient comprising:

(a) obtaining a sample of biological fluid from a patient; and

(b) detecting levels of PLA₂ in the sample;
10 wherein elevated levels of PLA₂ in the sample are indicative of cancer.

11. The method of claim 10 wherein the biological fluid is serum.

12. The method of claim 10 wherein PLA₂ is detected
15 by ELISA.

13. A method of monitoring progression, remission, response to therapy and stabilization of prostate, breast, ovarian or testicular cancer in a patient comprising measuring PLA₂ levels in biological fluid samples obtained from a patient
20 at selected times wherein an increase in the measured levels of PLA₂ over time is indicative of progressive cancer, a decrease in the measured levels of PLA₂ over time is indicative of remission or response to therapy of the cancer, and no change in the measured levels of PLA₂ over time is indicative
25 of stabilization of the cancer.

14. The method of claim 13 wherein the bodily fluid is serum.

- 20 -

15. The method of claim 13 wherein PLA_2 is measured by ELISA.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/03170

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : G01N 33/53, 53/567; C07K 16/00

US CL : 435/7.2, 7.21, 7.23, 7.91, 7.92; 436 546; 530/387.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/7.2, 7.21, 7.23, 7.91, 7.92; 436 546; 530/387.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, medline,

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98/05349 A1 (SMITHKLINE BEECHAM CORPORATION) 12 February 1998, see especially p.1, lines 10-11, p.2, lines 18-22 and Example 2.	13-15
X	YAMASHITA, S. et al. Elevation of Serum Group II Phospholipase A2 Levels in Patients with Advanced Cancer. Clinica Chimica Acta. 1994, Vol. 228, pages 91-99, see entire document.	13-15
X	ABE, T. et al. Group II Phospholipase A2 is Increased in Peritoneal and Pleural Effusions in Patients with Various Types of Cancer. Int. J. Cancer (Pred. Oncol.) 1997, Vol. 74, pages 245-250, see entire document.	13
-		---
Y		15

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*&* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

14 APRIL 1999

Date of mailing of the international search report

20 MAY 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

NANCY A. JOHNSON

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/03170

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	US 5,747,264 A (SCHMIDT, C.J. et al) 05 May 1998, see entire documnt.	13-15
A	YAMASHITA, S. et al. Overexpression of Group II Phospholipase A2 in Human Breast Cancer Tissues is Closely Associated With Their Malignant Potency. Cancer. 1994, Vol. 69, pages 1166-1170, see entire document.	1-15

Form PCT/ISA/210 (continuation of second sheet)(July 1992)★